Case Report

Exteriorized thrombosed axillo-bifemoral graft with empyema post coronary artery bypass grafting with axillo-femoral bypass

Anish Gupta^{1,*}, Bhavna Gupta²

¹Dept. of Cardiothoracic and Vascular Surgery, All India Institute of Medical Sciences, Rishikesh, Uttarakhand, India

²Dept. of Anesthesia, All India Institute of Medical Sciences, Rishikesh, Uttarakhand, India

ARTICLE INFO

Article history:

Received 15-04-2021

Accepted 23-04-2021

Available online 14-05-2021

Keywords:

Exteriorized Axillobifemoral graft

Coronary artery bypass surgery

ABSTRACT

Following its introduction in 1963, axillofemoral bypass grafts have improved, with patency rates comparable to aortofemoral and aortobifemoral grafts. Graft infection and thrombosis may worsen axillofemoral bypasses, rendering subsequent management challenging in surgically high-risk patients. We present an unusual case of an exteriorized thrombosed axillofemoral graft infection in a 65 years' male who recently underwent coronary artery bypass grafting with axillobifemoral grafting, presented in sepsis with empyema of chest and infected graft was exteriorised. He developed foul smelling discharge and fever. The patient was treated with antibiotics and infected graft was removed. The image highlights the social stigma and debilitating complications caused by these grafts in developing countries.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Case presentation

A 65 year old male presented to cardiac surgical outpatient unit with complaints of fever, chest pain and pus discharge from the previous chest tube insertion site. He had history of Coronary artery bypass grafting with axillo-femoral bypass about one month ago at some peripheral hospital. On examination, he had low grade fever, but his vitals were normal. His dacron graft has been exposed below the chest and has come out of the subcutaneous tunnel. It was grossly infected and non-pulsatile. On Doppler examination, it was found to be thrombosed. He was having foul smelling purulent discharge from the graft site as well as chest tube insertion site.

2. Description of the employed technique

Since patient was not having rest pain, he was admitted and started on intravenous antibiotics according to pus culture sensitivity. After stabilization, he was taken in the operative room, all three anastomotic sites were exposed and graft

was divided and the exteriorized graft was removed. The patient's general condition improved and sepsis subsided. He was discharged and was planned to be re-evaluated at a later stage for his peripheral vascular disease.

3. Discussion

Infection of prosthetic grafts complicate 0.5–3.5% of all patients with a mortality rate of up to 75% and is considered a horrific consequence of aorto-iliac revascularization procedures.^{1–3} The definitive treatment is graft excision followed by revascularization via extra-anatomical or in situ reconstruction. However, in clinical practice, care is adapted to the patient's comorbidities, and Samson's updated Szilagyi classification system of extra-cavitary vascular graft infection which correlates extent of infection with prognosis.^{3,4} When axillo-femoral graft infections occur in patients who have limited revascularization options and are unable to tolerate major re-operative procedures, complications arise. In such cases, rather than graft excision, the best treatment choice could be graft rescue or conservative antibiotic administration. A

* Corresponding author.

E-mail address: drash_06@yahoo.co.in (A. Gupta).

Fig. 1:

contaminated vascular graft that has degraded through adjacent structures necessitates graft removal in the form of overt septicemia. Skervin et al⁵ reported a axillo-femoral bypass graft transgressing the chest wall with sepsis related

to the patent graft. They test clamped the graft and after ensuring viability of limb, they explanted it under local anesthesia. They emphasized that explanting an infected extra-anatomical bypass graft does not need a concomitant revascularization procedure if the patient is at high risk of surgery.

Our report highlights the social stigma faced by the patients and debilitating complications that can be caused by these grafts especially in developing nations.

4. Source of Funding

None.

5. Conflict of Interest

None.

References

1. Mirzaie M, Schmitto JD, Tirilomis T, Fatehpur S, Liakopoulos OJ, Teucher N, et al. Surgical Management of Vascular Graft Infection in Severely Ill Patients by Partial Resection of the Infected Prosthesis. *Eur J Vasc Endovasc Surg.* 2007;33(5):610–3.
2. Garot M, Delannoy PY, Meybeck A, Bournet BS, Elia P, Escrivan T, et al. Intra-abdominal aortic graft infection: prognostic factors associated with in-hospital mortality. *BMC Infect Dis.* 2014;14(1):215.
3. Revest M, Camou F, Senneville E, Caillon J, Laurent F, Calvet B, et al. Medical treatment of prosthetic vascular graft infections: Review of the literature and proposals of a Working Group. *Int J Antimicrob Agents.* 2015;46(3):254–65.
4. Cappello E, Lorenzo MD, Cutillo E, Franco E. Axillofemoral bypass in elderly patients with local anesthesia: an alternative route to less risk. *BMC Surg.* 2013;13(1):6.
5. Skervin AL, Thomas RH, Sritharan K. An axillofemoral bypass graft transgressing the chest wall. *Clin Case Rep.* 2016;4(1):32–4.

Author biography

Anish Gupta, Assistant Professor

Bhavna Gupta, Assistant Professor

Cite this article: Gupta A, Gupta B. Exteriorized thrombosed axillo-bifemoral graft with empyema post coronary artery bypass grafting with axillo-femoral bypass. *IP J Surg Allied Sci* 2021;3(1):25-26.