Case Report

Ritual circumcision by inexperienced hand some time cause penile amputation

Dinesh Kumar Barolia^{1,*}, Arka Chatterjee², Gunjan Sharma², Ramesh Tanger²,
Arun Kumar Gupta²

¹Dept. of Paediatric Surgery, J.L.N. Medical College, Ajmer, Rajasthan, India

²Dept. of Paediatric Surgery, Sir Padampat Mother and Child Health Institute, S.M.S. Medical College, Jaipur, Rajasthan, India

ARTICLE INFO

Article history:

Received 16-08-2021

Accepted 20-08-2021

Available online 18-10-2021

Keywords:

Circumcision

Glans penis

Ritual circumcision

Penile amputation

ABSTRACT

Background: Ritual circumcision is a custom. But, when it is done by inexperienced person, it may cause painful situation for parents.

Case Presentation: We are presenting a case of ritual circumcision done by inexperienced person. Complete amputation of glans was occurred during circumcision. That was very painful situation for parents.

Conclusion: To avoid such complication circumcision should be done by experienced medical practitioner or training should be provided to the persons who are doing ritual circumcision.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Circumcision is the surgical procedure in which removal of the foreskin covering the glans penis. It is the one of the oldest surgical procedure practiced worldwide.¹ The term circumcision is derived from Latin word *circumcidere*, meaning "to cut around".² Reasons behind the circumcision are cultural, religious, medical or cosmetic. There are various technics commonly used to do circumcision. These entire technic have the main goal is to do circumcision with minimal bleeding and complications.³ The most common neonatal circumcision complication is bleeding, infection and irrational removal of foreskin.⁴ Amputation of glans is the most devastating complication in neonatal circumcision. We are reporting here a case penile amputation during circumcision, a drastic complication.

2. Case Report

A 45 days old male child born with full term normal vaginal institutional delivery was admitted at our institution with complaint of continuous bleeding after ritual circumcision. On clinical examination left radial pulse rate was 107 per minute, hypovolemic, feeble. Respiratory rate was 39 per minute. He was clinically pallor. On local examination the glans penis was complete separated from penile shaft. A bit of skin was holding the distal amputated glans and penile shaft otherwise circumferentially skin was incised. Baby was optimized first. Blood sample was sent for cross match, baseline investigations. Blood transfusion was done. Risk and prognosis was explained to parents and written consent was taken for surgical repair. Baby was taken for surgery after optimization. He was operated under general anaesthesia with endotracheal intubation. After painting and draping local region was cleaned again with povidine solution. A 6 french silicon foley's catheter was introduced through glans meatus and then introduced in penile shaft urethra. Foleys catheter balloon was inflated with 2 ml

* Corresponding author.

E-mail address: dbaroliarnt@gmail.com (D. K. Barolia).

normal saline. With this catheter stent glans was super imposed over amputated end of penile shaft in anatomical position and simple repair was done circumferentially with vicryl suture in interrupted pattern. Over this skin was repaired. Dorsal slitting of prepuce was done to observe the viability of glans during dressing and their after. Shukla dressing was done to prevent excessive movement of repaired tissue and catheter. First dressing was done after five days. The glans was viable and stitch line of skin was intact. Second dressing was done after five days of first dressing. Glans was intact and slight gaping was present at left lateral stitch line, otherwise normal. Skin was healed with secondary intention. There was no pus discharge or urine leak from wound site. After three days dressing was removed and Foleys catheter kept intact. The wound kept open. He was discharged on oral medication with Foleys catheter insitu. Foleys catheter was removed after six weeks of repair. After removal urine stream was straight and coming out from urethral meatus. There was no urethrocutaneous fistula. Six month follow up was uneventful.

Fig. 1: Complete transected penile shaft

3. Discussion

Circumcision in neonates or children is commonly done for either ritual or cultural reason or medical reason.^{5,6} It is commonly done in Muslims, Jews, Southeast Asia and Africa and in United States.^{6,7} In Jews community circumcision of male infant performed on live day eight. The Torah holy book of Jews state that it was the covenant between Abraham and God. This covenant confirmed by male infant circumcision. Muslims do circumcision for determination of relationship with God and purification.⁶

Fig. 2: Stenting of distal transected penis

Fig. 3: Stenting of both distal transected and proximal intact penis

Fig. 4: After repair of amputated penis

Fig. 5: Penile look after three months of repair

Circumcision used to perform from ancient time. Various complications were reported with time. The Complications after neonatal circumcision were observed by Sebastian O. Ekenze et al. in their study and reported in 2013. They found glanular adhesion, meatal stenosis, urethrocutaneous fistula, trapped penis, implantation dermoid, and glans amputation after circumcision.⁸ Penile amputation is really a traumatic to children and their parents. To decrease these complications with time various methods and apparatus were introduced in medical science. The techniques used for circumcision are Shield and clamp, Plastibell, Gomco, Zhenxi Rings, Tara Klamp, Smart Klamp, Shang Ring, PrePex, and Laser circumcision.⁹

Penile replantation was first reported by Ehrich WS (1929) in self-inflicted penile amputation by a man with radial saw.¹⁰ But successful replantation was reported in 1977. The replantation was done by micro neurovascular repair. First they repair urethra with stenting 8 french catheter.¹¹ We also repair first urethra over 6 french silicon catheter stent. After it corpora spongiosum is repaired and then after corpora cavernosa were repaired in anatomical position. Covering skin was approximated after refreshing the margin. Micro vascular anastomosis was not possible, due to some circumstances. Similar replantation was done by Bouassida Khairedine et al. (2012) in two cases and found good results.¹²

4. Conclusion

Circumcision is a procedure which performed commonly worldwide. Reasons behind circumcision are different, but it should be performed by experienced medical practitioner. And non-medical person doing ritual or cultural circumcision should be trained by medical professional to avoid complications.

5. Source of Funding

None.

6. Conflict of Interest

The authors declare no conflict of interest.

References

- Blank S, Brady M, Buerk E, Carlo W, Diekema D, Freedman A, et al. American Academy of Pediatrics, Task force on Circumcision: Male circumcision. *Pediatrics*. 2012;130:756–85.
- Lissauer T, Clayden G. Illustrated Textbook of Paediatrics. Fourth edition ed. Elsevier; 2011. p. 352–3.
- Gluckman GR, Stoller ML, Jacobs MM, Kogan BA. Newborn penile glans amputation during circumcision and successful reattachment. *J Urol*. 1995;153(3):778–9.
- Krill AJ, Palmer LS, Palmer JS. Complications of Circumcision. *Sci World J*. 2011;11:2458–68. doi:10.1100/2011/373829.
- American Academy of Pediatrics Task Force on Circumcision. Male circumcision. *Pediatrics*. 2012;130:756–85.
- WHO/UNAIDS. Male Circumcision: Global Trends and Determinants of Prevalence, Safety and Acceptability. World Health Organization; 2008.
- Owings M, Uddin S, Williams S. Health E-Stat “Trends in Circumcision for Male Newborns in U.S. Hospitals: 1979-2010”. National center for health statistics. Available from: https://www.cdc.gov/nchs/data/hestat/circumcision_2013/Circumcision_2013.htm.
- Ekenze SO, Ezomike UO. Complications of neonatal circumcision requiring surgical intervention in a developing country. *J Trop Pediatr*. 2013;59(4):292–299.
- Abdulwahab-Ahmed A, Mungadi IA. Techniques of male circumcision. *J Surg Tech Case Report*. 2013;5:1–7.
- Ehrich WS. Two unusual penile injuries. *J Urol*. 1929;21:239.
- Cohen BE, May JW, Daly JS, Young HH. Successful clinical replantation of an amputated penis by microneurovascular repair. Case report. *Plast Reconstr Surg*. 1977;59(2):276–80.
- Khairiddine B, Adnen H, Khaled BB, Adel S. Surgical reimplantation of penile glans amputation in children during circumcision. *Urol Ann*. 2014;6:85–7.

Author biography

Dinesh Kumar Barolia, Assistant Professor <https://orcid.org/0000-0003-0617-8435>

Arka Chaterjee, Senior Resident

Gunjan Sharma, Senior Resident

Ramesh Tanger, Associate Professor

Arun Kumar Gupta, Senior Professor

Cite this article: Barolia DK, Chaterjee A, Sharma G, Tanger R, Gupta AK. Ritual circumcision by inexperienced hand some time cause penile amputation. *IP J Surg Allied Sci* 2021;3(3):77-80.